



# Lecture 7

## Derivatives of exp and ln

### 7.1 Drawing derivatives of a graph

**Exercise 7.1.1.** Below on the left is the graph of  $f(x) = e^x$ .



Let me tell you the following fact: The derivative of  $e^x$  at  $x = 0$  is 1. (In fact, the value of  $e^x$  at  $x = 0$  is 1 also.)

- (a) Based on this, draw the derivative of  $e^x$  on the right.
- (b) How does your drawing compare to the graph of  $e^x$ ?

In fact, we have the following theorem:

**Theorem 7.1.2** (Derivative of  $e^x$ ). The derivative of  $e^x$  is itself. That is,

$$(e^x)' = e^x.$$

Put another way,

$$\frac{d}{dx}(e^x) = e^x.$$

How cool is that? There's a function that is its *own* derivative!

**Example 7.1.3.** Let's find the derivative of  $e^{3x}$ . We have

$$\frac{d}{dx}(e^{3x}) = \frac{d}{dx}(3x) \cdot \frac{d(e^x)}{dx}(3x) \tag{7.1.1}$$

$$= 3 \cdot e^{3x}. \tag{7.1.2}$$

We have used the chain rule in the first line. If you're confused by it, it may be worthwhile to write this out step-by-step. Let's let  $f(x) = e^x$  and  $g(x) = 3x$ . Then  $e^{3x} = f(g(x))$ . Thus

$$\frac{d}{dx}(e^{3x}) = \frac{d}{dx}f(g(x)) \tag{7.1.3}$$

$$= f'(g(x)) \cdot g'(x) \tag{7.1.4}$$

$$= f'(3x) \cdot g'(x). \tag{7.1.5}$$

(The second equality is due to the chain rule.) But we know that  $f'(x) = e^x$  by Theorem 7.1.2, and we know  $g'(x) = 3$  from previous lectures. Hence we can continue:

$$f'(3x) \cdot g'(x) = e^{3x} \cdot 3 = 3e^{3x}.$$

**Exercise 7.1.4.** Fix a real number  $B$ . Prove that the derivative of

$$f(x) = e^{Bx}$$

equals

$$Bf(x).$$

More generally, if you have another real number  $A$ , let

$$g(x) = Ae^{Bx}.$$

(For example, if you choose  $A = 3$  and  $B = 5$ , you would have  $3e^{5x}$ . The previous example is when  $A = 1$ .) Prove that

$$g'(x) = Bg(x).$$

**Application 7.1.5.** This kind of behavior is incredibly important for *modeling*. For example, how fast is a population growing? In ideal circumstances, the more individuals there are in a population, the faster we expect the population to grow. Better yet, we might expect that the rate of population growth is *proportional* to the population itself! (Note that “being proportional to” is a far more precise relationship than “the bigger the population, the faster the growth”.)

That’s exactly what Exercise 7.1.6 tells us about  $g(x) = Ae^{Bx}$ . We see that  $g'$  is proportional to  $g$  (with proportional constant  $B$ ). So for example,  $x$  could model time, while  $g(x)$  could model the population at time  $x$ .

By the way, why might  $g(x)$  be a *bad* model for population growth? For what kinds of situations might  $g(x)$  be a *good model*? In those situations, what might  $A$  and  $B$  represent?

**Exercise 7.1.6.** Find the derivative of  $f(x) = 5^x$ . Hints: Remember that  $5 = e^{\ln 5}$ , remember the basic rules for dealing with exponents, and use the chain rule.

**Exercise 7.1.7.** Your friend is excited about the idea that  $f(x)$  could equal  $f'(x)$  and looks for more examples that look like  $e^x$ . They try  $f(x) = 5^x$ , and are disappointed that  $f'(x)$  does not equal  $f(x)$ .

Is it possible to find any number  $k$ —other than  $e$ —so that if  $f(x) = k^x$ , then  $f'(x) = f(x)$ ?

**Remark 7.1.8.** Isn’t  $e$  special?

**Exercise 7.1.9.** Now that you know the derivative of  $g(x) = e^x$ , can you figure out the derivative of  $f(x) = \ln x$ ?

Hint: What is  $g \circ f$ ? What if you try computing  $(g \circ f)'$  using the chain rule, too?


## 7.2 Review of right inverses

Let  $f$  be a function. Here's a question: Given a value of  $f$ , can we always determine which  $x$  it came from?

**Example 7.2.1.** Here are some examples:

1. If  $f(x) = 3x$ , and if someone tells you that  $f$  takes the value 12, you know exactly where:  $x$  must equal 4. In fact, in general, if  $f$  takes value  $y$ , you know the original  $x$  is  $y/3$ .
2. If  $f(x) = 2^x$ , and if someone tells you that  $f$  takes the value 8, you know exactly where:  $x$  must equal 3. In fact, in general, if  $f$  takes value  $y$ , you know  $f$  does so at  $\log_2 y$ .
3. If  $f(x) = x^2$ , and if someone tells you that  $f$  takes the value 4, you *don't* know exactly where:  $x$  could equal 2 or -2. However, *if* you restrict yourself to looking only for positive values of  $x$ , then if  $f$  takes value  $y$ , you know that the original  $x$  is  $\sqrt{y}$ .

Below is a visual way to think about this process. Drawn is the graph of  $f$ . Given a value  $y$ , can you figure out which value of  $x$  satisfies the equation  $f(x) = y$ ? If so, that means that the coordinate  $x$  now becomes a function of  $y$ —you input  $y$ , and you output  $x$ —and we can call this function  $g$ .



**Warning 7.2.2.** While we were diligent about drawing  $g$  as a function of  $y$  before, from now on, we must now be comfortable realizing that letters are just letters, and we don't care if  $g$  takes inputs to be symbols that look like " $x$ ," or symbols that look like " $y$ "; that is,  $g$  will often be treated as a function of  $x$ , too.

**Definition 7.2.3.** Let  $f$  be a function. We say that a function  $g$  is a *left inverse* to  $f$  if

$$(g \circ f)(x) = x.$$

Put another way,  $g$  “remembers” the original value  $x$  that outputted  $f(x)$ .

We also say that  $f$  is a *right inverse* to  $g$ . Put another way,  $f$  “knows” that if  $g(\text{something}) = x$ , then  $\text{something} = f(x)$ .

### 7.3 Derivatives of right inverses

It turns out that if we know the derivatives of a function  $g$ , then—if  $g$  has a right inverse  $f$ —we can figure out the derivatives of the right inverse  $f$ .

**Lemma 7.3.1.** Let  $g$  be a function, and suppose that  $f$  is a right inverse to  $g$ , defined on some open interval containing  $x$ . Suppose also that  $g$  is differentiable at  $f(x)$ , and that  $g'(f(x)) \neq 0$ . Then

$$f'(x) = \frac{1}{g'(f(x))}. \quad (7.3.1)$$

That is, the derivative of  $f$  at  $x$  is computed by dividing 1 by the derivative of  $g$  at  $f(x)$ .

*Proof.* Let’s look at the following string of equalities:

$$1 = (x)' \quad (7.3.2)$$

$$= (g \circ f)'. \quad (7.3.3)$$

The first equality is our knowledge of the derivative of the function  $x$ . The next equality is using the hypothesis that  $g$  is a right inverse to  $f$ , so that  $f \circ g = x$ .

In total, what this string of equalities says is that the function on the righthand side is equal to the (constant!) function on the lefthand side. So let’s evaluate at some point  $x$ . We have

$$1 = (g \circ f)'(x) = g'(f(x)) \cdot f'(x).$$

We can divide both sides by  $g'(f(x))$  so long as this number isn’t zero; so we find:

$$\frac{1}{g'(f(x))} = f'(x) \quad \text{when } g'(f(x)) \neq 0.$$

This is what we wanted. □

**Example 7.3.2.**  $f = \ln(x)$  is a right inverse to  $g(x) = e^x$ . This is because

$$g \circ f(x) = e^{\ln x} = x.$$

We know the derivative of  $g(x) = e^x$ , so we can use the lemma to find the derivative of  $f(x) = \ln(x)$ ! Let's try:

$$(\ln(x))' = f'(x) \tag{7.3.4}$$

$$= \frac{1}{g'(f(x))} \tag{7.3.5}$$

$$= \frac{1}{e^{f(x)}} \tag{7.3.6}$$

$$= \frac{1}{e^{\ln(x)}} \tag{7.3.7}$$

$$= \frac{1}{x}. \tag{7.3.8}$$

. The first equality is by definition of  $f$ . The next equality is using Lemma 7.3.1. The rest is just plugging in our knowledge of  $g'$  and  $\ln$ .

## 7.4 The derivative of natural log

The example from the last page is important, so let's record this as a theorem. (You will be expected to know this:)

**Theorem 7.4.1** (The derivative of  $\ln$ ). The derivative of  $\ln$  is “one over  $x$ .” That is,

$$\frac{d}{dx} \ln(x) = \frac{1}{x}.$$

## 7.5 For next lecture

You should be comfortable finding derivatives of functions involves  $e^x$  and  $\ln$ . For example, you should be able to find  $f'$  for each of the following functions  $f$ :

(a)  $f(x) = e^x$

(b)  $f(x) = e^{3x}$

(c)  $f(x) = e^{3x+2}$

(d)  $f(x) = 3e^x$

(e)  $f(x) = 5^x$

(f)  $f(x) = 5^{3x}$

(g)  $f(x) = \ln(x)$

(h)  $f(x) = \ln(3x)$

(i)  $f(x) = \ln(x + 3)$