

# Lecture 10

## Second derivatives, concavity, and minima/maxima

### 10.1 Second derivatives

Today, we will practice taking “second derivatives,” and knowing when they are positive or negative.

**Definition 10.1.1.** The second derivative of  $f$  is *the derivative of the derivative*<sup>1</sup> of  $f$ . We denote the second derivative by

$$f'', \quad \text{or} \quad \frac{d}{dx}\left(\frac{d}{dx}f\right), \quad \text{or} \quad \frac{d^2}{dx^2}f, \quad \text{or} \quad \frac{d^2f}{dx^2}. \quad (10.1.1)$$

**Example 10.1.2.** Let  $f(x) = 3x^2 + x - 7$ . Then the (first) derivative of  $f$  is

$$f'(x) = 6x + 1.$$

If we take the derivative of  $f'(x)$ , we end up with the second derivative of  $f$ :

$$f''(x) = 6.$$

**Example 10.1.3.** Here are more examples of functions and their second derivatives. You should verify these examples:

- If  $f(x) = \sin(x)$ , then  $f''(x) = -\sin(x)$ .

---

<sup>1</sup>Yes, there are two appearances of the word “derivative”; this is not a typo.

10LECTURE 10. SECOND DERIVATIVES, CONCAVITY, AND MINIMA/MAXIMA

- If  $f(x) = e^x$ , then  $f''(x) = e^x$ .
- If  $f(x) = e^{5x}$ , then  $f''(x) = 25e^{5x}$ .
- If  $f(x) = x^3 - 5x^2$ , then  $f''(x) = 6x - 10$ .

**Example 10.1.4.** Let's find the second derivative of  $f(x) = \ln(x)$ . As defined above, we just need to take the derivative twice. Let's take the first derivative:

$$f'(x) = \frac{1}{x}.$$

(This is something we learned in class.) Now let's take another derivative—for example, by using the quotient rule—to find

$$f''(x) = \frac{0 \cdot x - 1 \cdot 1}{x^2} = -\frac{1}{x^2}.$$

That is, the second derivative of  $\ln x$  is  $-1/(x^2)$ .

**If you know how to take derivatives, you know how to take second derivatives.** So you see how our skills are building on each other—make sure you practice taking derivatives!

**Example 10.1.5.** Let  $f(x) = x^2 - 2$ . *Where is the second derivative positive?*

Let's find the second derivative. We see that

$$f'(x) = 2x$$

so

$$f''(x) = 2.$$

So the second derivative is always 2, meaning the second derivative is positive *everywhere*.

**Example 10.1.6.** Let  $f(x) = x^3 - 3x^2 + 3$ . *Where is the second derivative positive?*

Let's find the second derivative. We see that

$$f'(x) = 3x^2 - 6x$$


so, taking the derivative of  $f'(x)$ , we find:

$$f''(x) = 6x - 6.$$

So the second derivative is positive when  $6x - 6$  is positive. This happens exactly when  $6x > 6$ —that is, when  $x > 1$ .

As a bonus: The second derivative is negative when  $6x < 6$ —that is, when  $x < 1$ .

Below is a graph of  $f(x)$ , and I have shaded in **bold** the part of the graph where the second derivative is positive:



**Example 10.1.7.** Let  $f(x) = x^4 - 24x^2 + 50$ . Where is the second derivative positive?

Let's find the second derivative. We see that

$$f'(x) = 4x^3 - 48x$$

so, taking the derivative of  $f'(x)$ , we find:

$$f''(x) = 12x^2 - 48.$$


So the second derivative is positive when  $12x^2 - 48$  is positive. This happens exactly when  $12x^2 > 48$ —that is, when  $x^2 > 4$ . But  $x^2 > 4$  exactly when  $x < -2$  or  $x > 2$ .

As a bonus: The second derivative is negative when  $x^2 < 4$ —that is, when  $x$  is between  $-2$  and  $2$ .

Below is a graph of  $f(x)$ , and I have shaded in **bold** the part of the graph where

12LECTURE 10. SECOND DERIVATIVES, CONCAVITY, AND MINIMA/MAXIMA

the second derivative is positive:



**Example 10.1.8.** Let  $f(x) = 3 \sin(x)$ . Where is the second derivative positive?

Let's find the second derivative. We see that

$$f'(x) = 3 \cos(x)$$


so, taking the derivative of  $f'(x)$ , we find:

$$f''(x) = -3 \sin(x)$$


So the second derivative is positive when  $-3 \sin(x)$  is positive. This happens exactly when  $\sin(x)$  is negative. And based on our trigonometry knowledge from precalculus, we know that this happens when

- $x$  is between  $\pi$  and  $2\pi$ ,
- $x$  is between  $3\pi$  and  $4\pi$ ,
- $x$  is between  $-\pi$  and  $0$ ,
- $x$  is between  $-3\pi$  and  $-\pi$ ,
- ....

Below is a graph of  $f(x)$ . I have shaded in **bold** the part of the graph where the second derivative is positive:




For next class, I expect you to be able to do the following: For each of the functions  $f(x)$  below, (i) State *where* the function has a *positive* second derivative, and (ii) Shade in **bold** where the graph of the function has positive second derivative. (You will be provided the graph of  $f(x)$ .)




(a)  $f(x) = -x^4 + 24x^2 - 50$ .


14LECTURE 10. SECOND DERIVATIVES, CONCAVITY, AND MINIMA/MAXIMA



(b)  $f(x) = x^2$ .




(c)  $f(x) = \cos(x)$ .




(d)  $f(x) = \tan(x)$ .

You have seen examples of graphs with positive second derivative. Here are some examples, with the positive-second-derivative regions shaded in *bold*:


1.  $f(x) = x^3 - 3x^2 + 3$ :




3.  $f(x) = x^4 - 24x^2 + 50$ :




2.  $f(x) = 3 \sin(x)$ :



4.  $f(x) = e^x$ :



5.  $f(x) = \tan(x)$ :



## 10.2 Concavity

The point I want to make with these pictures is that *the value of the second derivative gives us some idea of what the graph looks like.* (Though not a complete picture.)

Intuition: On the regions where the second derivative is positive, the graph of  $f$  looks like a *portion* of an “upright bowl.” Some students have described this as “opening upward” as well.

Conversely, when the second derivative is negative, the graph of  $f$  looks like a portion of an “upside-down bowl.” But we have technical names, too. From now on, you are expected to know the following terminology:


**Definition 10.2.1** (Concavity). We say that  $f$  is *concave up* at  $x$  if  $f''(x) > 0$ . We say that  $f$  is *concave down* at  $x$  if  $f''(x) < 0$ .

## 10.3 Inflection points


**Definition 10.3.1.** If  $f''(x) = 0$ , and the concavity of  $f$  *changes* at  $x$ , we say that  $x$  is an *inflection point*.

**Example 10.3.2.** Here are some examples of functions and their graphs, with their inflection points labeled.

1.  $f(x) = x^3 - 3x^2 + 3$ :




2.  $f(x) = 3 \sin(x)$ :






3.  $f(x) = x^4 - 24x^2 + 50$ :




4.  $f(x) = e^x$ :




(No inflection points.)

5.  $f(x) = \tan(x)$ :



6.  $f(x) = x^4$ :

(No inflection points, even though  $f''(x) = 0$  at  $x = 0$ .)

**Expectation 10.3.3.** Based on looking at a graph, you are expected to be able to identify inflection points—an inflection point is a place at which a function switches concavity (from up to down, or from down to up).

**Exercise 10.3.4.** Now we're going to try to learn something about a function by knowing its derivative and second derivative. You can hunt for examples on the previous pages of this packet. Or, you can try understanding  $f(x) = x^2$  and  $f(x) = -x^2$ .

1. Can you find an example of a function  $f$ , and a point  $x$ , where  $f'(x) = 0$  and  $f$  is concave up at  $x$ ? What does the function  $f$  look like near  $x$ ? How does the value of  $f$  at  $x$  compare to the value of  $f$  at nearby points?
2. Can you find an example of a function  $f$ , and a point  $x$ , where  $f'(x) = 0$  and  $f$  is concave *down* at  $x$ ? What does the function  $f$  look like near  $x$ ? How does the value of  $f$  at  $x$  compare to the value of  $f$  at nearby points?

## 10.4 For next time

You should be able to tell me the second derivatives of the following functions:

- (a)  $x^3 - 3x^2 + x$
- (b)  $4x^2 + 3x - 2$
- (c)  $e^{7x}$
- (d)  $\sin(x)$

You should also be able to tell me where the following functions are concave up:

- (a)  $x^3 - 3x^2 + x$
- (b)  $4x^2 + 3x - 2$
- (c)  $e^{7x}$